

1. (a) Show that if  $u \in h_p$  for  $p \geq 1$ , then  $|u(z)| \leq \left(\frac{1+|z|}{1-|z|}\right)^{1/p} \|u\|_p$ .
- (b) Show that there is a universal constant  $C_{1,p}$  so that if  $f \in H^p$  for  $p \geq 1$ , then
 
$$|f'(z)| \leq C_{1,p}(1-|z|)^{-1-1/p} \|f\|_p.$$

**Hint:** consider  $g(z) = \frac{f(z)-f(z_0)}{z-z_0}$ .

- (c) Generalize to find a similar estimate for the  $n$ th derivative of  $f \in H^p$ .
2. (a) Show that if  $K \subset \mathbb{C}$  is compact and  $v : K \rightarrow [-\infty, +\infty)$  is u.s.c., then there is a decreasing sequence  $u_n$  of continuous functions decreasing to  $v$ .
- (b) Show that if  $v$  is subharmonic on a connected open set  $\Omega$  and is not constantly  $-\infty$ , then for every  $\overline{D_r(z_0)} \subset \Omega$ ,

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} v(z_0 + re^{it}) dt > -\infty.$$

- (c) Show that if  $v_1$  and  $v_2$  are subharmonic on  $\Omega$ , then so is  $\max\{v_1, v_2\}$ .
- (d) Prove that if  $v$  is  $C^2$  and  $\Delta v \geq 0$  on  $\Omega$ , then  $v$  is subharmonic.
3. (a) Rewrite the formula for the conjugate function of  $u = f * P$  for  $f \in L^1$  as follows:

$$\tilde{u}(re^{i\theta}) = \frac{1}{2\pi} \int_0^\pi \left( \frac{4r \sin^2(t/2)}{(1-r)^2 + 4r \sin^2(t/2)} \right) \frac{f(\theta-t) - f(\theta+t)}{\tan(t/2)} dt.$$

Hence show that if  $\frac{1}{2\pi} \int_0^\pi \left| \frac{f(\theta-t) - f(\theta+t)}{\tan(t/2)} \right| dt < \infty$ , then  $\tilde{u}$  has a radial limit  $\lim_{r \rightarrow 1^-} \tilde{u}(re^{i\theta})$ .

- (b) Hence show that if  $f$  is  $C^1$  on an arc  $I \subset \mathbb{T}$ , then for every compact arc  $K \subset I$ ,  $\tilde{u}(re^{i\theta})$  converges to  $\frac{1}{2\pi} \int_0^\pi \frac{f(\theta-t) - f(\theta+t)}{\tan(t/2)} dt$  uniformly as  $z \in \mathbb{D}$  approaches  $e^{i\theta} \in K$ .
- (c) In particular, show that  $\tilde{u}$  extends to be continuous on  $\mathbb{D} \cup I$ .
4. Let  $K$  be a compact subset of  $\mathbb{T}$  of Lebesgue measure zero.

(a) Find a positive function  $f \in L^1$  such that  $f$  is  $C^\infty$  on  $\mathbb{T} \setminus K$  and  $\lim_{t \rightarrow t_0} f(t) = +\infty$  for every  $t_0 \in K$ .

(b) Define  $u = f * P$  and let  $\tilde{u}$  be its harmonic conjugate. Set  $g(z) = \frac{u + i\tilde{u}}{1 + u + i\tilde{u}}$ .

Prove that  $g \in A(\mathbb{D})$  (i.e.  $g$  extends to be continuous on the closed disc  $\overline{\mathbb{D}}$ ), that  $g|_K = 1$  and  $|g| < 1$  on  $\overline{\mathbb{D}} \setminus K$ .

(c) If  $\mu \in M(\mathbb{T})$  is of analytic type, evaluate  $\lim_{k \rightarrow \infty} \int g^k d\mu$  in two ways. Hence provide another proof of the F. and M. Riesz Theorem.

5. (a) Show that if  $f \in H^p$  and  $w$  is an analytic function from  $\mathbb{D}$  into itself, then  $f(w(z))$  belongs to  $H^p$ . **Hint:** use the least harmonic majorant.
- (b) Show that  $f(z) = (1-z)^{-1}$  is in  $H^p$  for  $0 < p < 1$ , but not  $p = 1$ .
- (c) Show that if  $f$  is analytic on  $\mathbb{D}$  and  $\operatorname{Re} f(z) > 0$ , then  $f \in H^p$  for  $0 < p < 1$ .
- (d) Generalize to analytic functions with values in a sector  $S = \{z : \alpha < \arg f(z) < \beta\}$ .